

EPA published the final [human health toxicity assessment for PFOA](#) in April 2024.

A toxicity assessment summarizes the potential health effects associated with exposure to a particular chemical and identifies the dose levels at which the health effects may occur in order to calculate toxicity values.

The PFOA toxicity assessment identified adverse health effects associated with PFOA exposure using a robust systematic review process based on [EPA peer-reviewed human health risk assessment methodology](#).

Systematic review is a structured and documented process for transparent literature review using explicit, pre-specified scientific methods to identify, select, assess, and summarize the findings across relevant studies.

Systematic review promotes use of the best available science and reduces bias.

The EPA followed its peer-reviewed human health risk assessment methodology and applicable guidance documents for all steps of the toxicity assessment including hazard identification, cancer classification, and toxicity value development (e.g., [USEPA, 2002](#); [USEPA, 2005](#); [USEPA, 2012](#); [USEPA, 2022](#)). The PFOA toxicity assessment incorporated [expert scientific recommendations received from peer review](#) and [feedback from the public comment period](#).

Health Effects Identified for PFOA

The EPA's systematic review of over [780 human and animal health studies](#) demonstrated PFOA exposure elicits adverse noncancer and cancer health effects (see table below). Consistent with [EPA's Guidelines for Carcinogen Risk Assessment](#), the EPA concluded that PFOA is Likely to Be Carcinogenic to Humans via the oral route of exposure.

	Immune	Developmental	Cardiovascular	Liver	Cancer
Effects observed in human studies	↓ Vaccine response in children	↓ Infant birth weight	↑ Serum lipids (total cholesterol and LDL) ↑ Blood pressure in adults	↑ Serum liver enzymes (ALT) indicative of liver damage in adults	Kidney and testicular cancer in adults
Effects observed in animal studies	↓ Immune response	↓ Pup survival ↓ Fetal and pup body weight Delayed developmental timing	Change in serum lipids	↑ Liver cell death and serum liver enzymes (ALT) indicative of liver damage	Testicular, pancreatic, and liver tumors

Toxicity Values for PFOA

Based on the effects described above, toxicity values were calculated for PFOA – a cancer slope factor (CSF) and a reference dose (RfD) – in line with EPA peer-reviewed human health risk assessment methodology and applicable guidance documents.

- A [CSF](#) is an upper bound, approximating a 95% confidence limit, on the increased cancer risk from a lifetime exposure to an agent. The CSF is the change in risk per unit dose.
- A [RfD](#) is an estimate (with uncertainty spanning perhaps an order of magnitude) of a daily oral exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious effects during a lifetime. Uncertainty in the data is accounted for by including uncertainty factors in the RfD to protect public health.

	Critical Effect	Study	Population	Toxicity Value
Cancer Slope Factor	Kidney cancer	Shearer, 2021 (supported by Viera, 2013; see Sec. 4.2.2.1)	55- to 74-year-old adults	0.0293 (ng/kg/day) ⁻¹
Reference Dose (see page 2)	↓ Tetanus and diphtheria vaccine response	Budtz-Jørgensen, 2018	5- to 7-year-old children	3×10 ⁻⁸ mg/kg/day
	↓ Birth weight	Wikström, 2020	Infants	
	↑ Serum total cholesterol	Dong, 2019	20- to 80-year-old adults	